



We create chemistry

GOLIATH® GEL

Reg. No. L 5763; N-AR 0873

Act No. 36 of 1947

Fipronil (phenyl pyrazole) 0,5 g/kg

WARNINGS

- Handle with care.
- Harmful if swallowed.
- Store in the original box in a cool place away from food and animal feed.
- Keep out of reach of children, uninformed people and animals.

PRECAUTIONS

- Prevent skin and eye contact. In case of accidental skin or eye contact wash with soap and water and rinse eyes thoroughly with clean water.
- Only apply out of the reach of children.
- Do not eat, drink or smoke whilst applying.
- Wash hands after application.
- Do not contaminate food processing surfaces with the gel. If GOLIATH® GEL contaminates non-target surfaces during application, remove the gel with a damp paper towel and throw away in a refuse bin. Clean area thoroughly.
- When empty, dispose of the cartridges safely.
- Do not attempt to open or re-fill the cartridge.
- Do not re-use the cartridge for any other purpose.
- Prevent contamination of food, feed, drinking water and eating utensils.

DIRECTIONS FOR USE

USE ONLY AS DIRECTED

General directions

GOLIATH® GEL is effective against the nymphs and adults of the German cockroach (*Blattella germanica*) and the American cockroach (*Periplaneta americana*).

GOLIATH® GEL may be used to control cockroaches in:

- Domestic premises (including kitchens).
- Food handling areas (i.e. for food processing, storage and preparation) such as food manufacturing premises, commercial kitchens, restaurants, food stores, warehouses, retail outlets etc.
- Public buildings such as hotels, hospitals, prisons, schools, churches, theatres etc.
- Commercial and industrial premises such as factories, shops, workshops, aircraft, vehicles, railway stock and ships.

Application

GOLIATH® GEL is most effective when placed within or close to cockroach harbourages, foraging and feeding areas. These are typically areas that are dark, warm, damp and remain undisturbed e.g. behind refrigerators and in electrical equipment. Locate such areas by appropriate techniques, such as inspection or by trapping with sticky traps. Search for empty egg cases, cockroach faeces and cast skins. For the best results the area to be treated should be cleaned and tidied before treatment, paying particular attention to the removal of alternative food sources.

GOLIATH® GEL cartridges are designed to be used with the BaitGun® applicator. Lock the GOLIATH® GEL cartridge into the BaitGun® and attach the appropriate application tip to the cartridge. Refer to the manufacturer's instructions for directions on the use of the BaitGun®.

GOLIATH® GEL should be applied throughout the infested area as small spots. Spots of gel should be applied in dark out-of-sight locations in cracks, crevices, behind cupboards and shelves, under kitchen appliances, in electrical control boxes, voids and ducting and under bathroom fixtures etc. Do not apply GOLIATH® GEL where it will become submersed or likely to be removed by routine cleaning. Do not spray other insecticides on or around GOLIATH® GEL spots or place spots on recently treated surfaces, as this will contaminate the GOLIATH® GEL and may discourage cockroaches from feeding on it.

GOLIATH® GEL is very attractive to cockroaches and feeding begins soon after placement of the gel. Typically cockroaches will die a few hours after a single feed on GOLIATH® GEL. In infested premises, dead cockroaches will normally be seen within 24 hours of treatment.

Treated areas should be re-inspected after 1 - 2 weeks. Where initial infestation was severe a second GOLIATH® GEL application may be required if the first treatment has been entirely consumed and live cockroaches are still present.

Rate of application

In cases of severe infestation and in areas that are particularly dirty or cluttered or where alternative sources of food cannot be entirely eliminated, use the higher rate of application.

Normal rate

German cockroach: One spot of 0,03 g gel per square metre.
 American cockroach: Two spots of 0,03 g gel per square metre.

High rate


German cockroach: Two spots of 0,03 g gel per square metre.
 American cockroach: Three spots of 0,03 g gel per square metre.

A spot of 0,03 g is approximately 3 - 4 mm in diameter.

The right number of spots must be placed per square metre of floor space (i.e. in a building of 100 m² floor space a total of 100 spots must be placed for the control of German cockroaches at the normal rate). This does not, however, mean that the spots must be placed evenly throughout the treated premises. Spots can be concentrated at cockroach harbourages and in other areas mentioned above. Cockroaches can often be found high up against ceilings and against walls if suitable shelter etc. is available there.

BASF SA (PTY) LTD acknowledges the following Registered Trademark:

BaitGun® is the Trademark of EFD Inc., U.S.A



GOLIATH® GEL

Reg. Nr: L 5763

Wet Nr 36 van 1947

Fipronil (feniel pirasool) 0,5 g/kg

WAARSKUWINGS

- Hanteer versigtig.
- Skadelik indien ingesluk word.
- Berg in die oorspronklike houër in 'n koel plek weg van voedsel en dierevoer.
- Hou buite bereik van kinders, oningeligte persone en diere.

VOORSORGMATREËLS

- Voorkom aanraking met vel en oë. In geval van toevallige aanraking met vel of oë was met seep en water en spoel oë deeglik met skoon water.
- Plaas die lokaas slegs buite die bereik van kinders uit.
- Moenie eet, drink of rook tydens toediening nie.
- Was hande na toediening.
- Moenie voedselverwerkingsoppervlakke met die gel besmet nie. As nie-teiken areas met **GOLIATH® GEL** besmet word tydens toediening, verwyder die gel met 'n klam papierhanddoek en gooi dit weg in 'n vullisdrom. Maak die area deeglik skoon.
- Wanneer leeg, raak op 'n veilige manier ontslae van die buis.
- Moenie probeer om die buis oop te maak of om dit te hervul nie.
- Moet buis vir geen ander doel hergebruik nie.
- Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS VOORGESKRYF

Algemene aanwysings

GOLIATH® GEL beheer die nimfe en volwassenes van die duitse kakkerlak (*Blattella germanica*) en die Amerikaanse kakkerlak (*Periplaneta americana*).

GOLIATH® GEL kan gebruik word vir die beheer van kakkerlakke in:

- Huishoudelike persele (insluitende kombuise).
- Plekke waar voedsel hanteer word (m.a.w. waar voedsel verwerk, geberg of voorberei word) soos in voedselverwerkingsaanlegte, kommersiële kombuise, restaurante, voedselstore, pakhuis, kleinhandelwinkels ens.
- Openbare geboue soos hotelle, hospitale, gevangenis, skole, kerke, teaters ens.
- Kommersiële en industriële persele soos fabriek, winkels, werkwinkels, vliegtuie, voertuie, spoorwegtrokke en skepe.

Toediening

GOLIATH® GEL is mees effektief wanneer dit in of naby kakkerlakskuilplekke en voedingsareas geplaas word. Hierdie areas is tipies donker, warm en vogtige plekke wat nie versteur word nie bv. agter yskaste en binne elektriese toebehore. Sulke plekke moet geïdentifiseer word deur deeglike inspeksie of die gebruik van taai lokvalle. Soek vir leë eiersakke, kakkerlakmis en vervellings. Vir die beste resultate moet die area wat behandel gaan word vooraf skoongemaak word. Veral alternatiewe voedselbronne moet verkieslik verwyder word.

Die **GOLIATH® GEL** buis is ontwerp om saam met die BaitGun® aanwender gebruik te word. Sit die **GOLIATH® GEL** buis in die BaitGun® aanwender en voeg die toepaslike aanwendingspunt aan die buis vas. Lees die vervaardiger se instruksies aangaande die gebruik van die Bait Gun®.

GOLIATH® GEL moet dwarsdeur die besmette perseel in klein kolletjies geplaas word. Kollie gel moet buite sig in donker plekke soos krate en gate, agter kaste en rakke, onder kombuistoehore, in elektriesekakelborde, in en om elektrieseleidings, onder badkamertoebehore ens. aangewend word. Moenie **GOLIATH® GEL** aanwend waar dit onder water gedompel kan word, of waar dit deur gewone skoonmaakprosesse verwyder kan word nie. Moenie ander insekdoders om of oor die **GOLIATH® GEL** kollie spuit nie of kollie op oppervlakke wat onlangs met insekdoders behandel is plaas nie. Dit sal die **GOLIATH® GEL** besmet en mag kakkerlakke ontmoedig om daaraan te vreet.

GOLIATH® GEL is baie aanloklik vir kakkerlakke en voeding op die lokaas begin baie gou na plasing van die lokaas. Gewoonlik vrek kakkerlakke binne enkele ure nadat hulle **GOLIATH® GEL** gevreet het. In besmette persele sal dooie kakkerlakke gewoonlik 24 uur na toediening sigbaar wees.

Behandelde areas moet na 1 - 2 weke weer geïnspekteer word. Waar die besmetting erg was, mag 'n tweede toediening van **GOLIATH® GEL** nodig wees indien al die vorige gel opgevrete is, en lewendige kakkerlakke nog steeds teenwoordig is.

Toedieningshoeveelheid

In gevalle van swaar besmetting of in areas wat baie vuil is of waar alternatiewe bronne van voedsel nie uitgesluit kan word nie, moet die hoër toedieningshoeveelheid gebruik word.

Normale toedieningshoeveelheid

Duitse kakkerlak: Een kol van 0,03 g gel per vierkante meter.

Amerikaanse kakkerlak: Twee kolle van 0,03 g gel per vierkante meter.

Hoër toedieningshoeveelheid

Duitse kakkerlak: Twee kolle van 0,03 g gel per vierkante meter.

Amerikaanse kakkerlak: Drie kolle van 0,03 g gel per vierkante meter.

'n Kol van 0,03 g is omtrent 3 - 4 mm in deursnee.

Die regte hoeveelheid kollie moet per vierkante meter vloerspasie geplaas word (m.a.w. in 'n gebou van 100 m² vloerspasie moet 100 kollie vir Duitse kakkerlakbeheer teen die normale toedieningshoeveelheid geplaas word). Dit beteken nie noodwendig dat kollie eweredig deur die perseel geplaas moet word nie. Kollie kan gekonsentreer word waar die kakkerlakke skuiling soek en in plekke soos hierbo beskryf. Kakkerlakke kan baie keer ook hoog op teen plafonne en mure gevind word as geskikte skuiling ens. op sulke plekke beskikbaar is.

BASF SA (PTY) LTD erken die volgende Handelsmerk:

BaitGun® is die Geregistreerde Handelsmerk van EFD Inc., U.S.A.